

**The following resources related to this article are available online at
www.sciencemag.org (this information is current as of November 15, 2009):**

Updated information and services, including high-resolution figures, can be found in the online version of this article at:

<http://www.sciencemag.org/cgi/content/full/303/5657/471b>

A list of selected additional articles on the Science Web sites **related to this article** can be found at:

<http://www.sciencemag.org/cgi/content/full/303/5657/471b#related-content>

This article **cites 6 articles**, 1 of which can be accessed for free:

<http://www.sciencemag.org/cgi/content/full/303/5657/471b#otherarticles>

This article appears in the following **subject collections**:

Geochemistry, Geophysics

http://www.sciencemag.org/cgi/collection/geochem_phys

Technical Comments

http://www.sciencemag.org/cgi/collection/tech_comment

Information about obtaining **reprints** of this article or about obtaining **permission to reproduce this article** in whole or in part can be found at:

<http://www.sciencemag.org/about/permissions.dtl>

Comment on "Impact Ejecta Layer from the Mid-Devonian: Possible Connection to Global Mass Extinctions"

The interesting report by Ellwood *et al.* (1) demonstrated the applicability of magnetic susceptibility field methods to the identification of distal impact deposits. However, the evidence that they have presented for a bolide impact during the Middle Devonian, immediately preceding the Eifelian-Givetian (E-G) stage boundary at ~380 million years ago (Ma), requires further confirmation and, in particular, better evidence for shock metamorphism. The images identified by Ellwood *et al.* (1) as shocked quartz grains are not convincing, and the orientation measurements suffer from an insufficient number of observations. The possible presence of an extraterrestrial component also needs confirmation: As and V anomalies are certainly not indicative of a cosmic component, and Ir data, which would be more characteristic, are absent.

Even if there were strong evidence for a mid-Devonian impact, the interpretation presented by Ellwood *et al.* (1) would be too far-reaching in one key respect. In the title of the report, Ellwood *et al.* announce a possible cause-and-effect relation between the putative bolide impact and a global mass extinc-

tion. This extinction in question is the late Eifelian Kacák/*otomari* event (2–4). Ellwood *et al.* claim that this biospheric perturbation "may represent the extinction of as many as 40% of all marine animal genera" (1). They cite the work of Sepkoski (5) as their only reference for this claim, but the 40% figure is not derived from this work. The crucial figure 6 from Sepkoski (5) illustrates the extinction pattern of only "well-preserved" marine genera during the Devonian substages, and the E-G passage is marked by demise level of approximately 15% (Fig. 1). More significant, however, is that this so-called mass extinction of the late Eifelian is at the sixth position among the Devonian intervals in a relative ranking, and has only half the magnitude of the peak that marks the late Frasnian event, a true mass extinction. Indeed, Walliser (2) characterized the impact-related Kacák/*otomari* biotic crisis as a minor two-step event that constituted at most a third-order extinction limited primarily to pelagic biota. Its main cause is seen to be the anoxic conditions recorded in widespread, long-lasting (perhaps 1 million years in duration) black-shale deposition during a pulse of ma-

rine transgression (2–4). The sulfide-forming and redox-sensitive trace metals, reported by Ellwood *et al.* (1) as evidence for an impact, may be alternatively thought as a geochemical signature of oxygen deficiency only.

Altogether, the term "global Kacák/*otomari* mass extinction" is an original proposal of Ellwood *et al.* (1), and an overestimate of the actual biotic changes at this stratigraphic horizon. Similarly negligible ecosystem collapse, at least in an immediate feedback, is established for the well-dated mid-Frasnian Alamo impact event in Nevada (6). The insistent search for causal link between an extraterrestrial impact of unknown magnitude and an obligatorily major extinction represents a circular argument [see, for example, (7, 8)], and such a link has not been demonstrated in this case.

Grzegorz Racki

Department of Earth Sciences
University of Silesia
PL-41-200
Sosnowiec, Poland
E-mail: racki@us.edu.pl

Christian Koeberl

Department of Geological Sciences
University of Vienna
Althanstrasse 14
A-1090 Vienna, Austria.
E-mail: christian.koeberl@univie.ac.at

References and Notes

1. B. B. Ellwood, S. L. Benoist, A. El Hassani, C. Wheeler, R. E. Crick, *Science* **300**, 1734 (2003).
2. O. H. Walliser, in *Global Events and Event Stratigraphy in the Phanerozoic*, O. H. Walliser, Ed. (Springer-Verlag, Berlin, 1996), pp. 225–250.
3. R. Schöne, *Göttinger. Arb. Geol. Paläont.* **70**, 1 (1997).
4. M. R. House, *Palaeogeogr. Palaeoclimatol. Palaeoecol.* **181**, 5 (2002).
5. J. J. Sepkoski Jr., in *Global Events and Event Stratigraphy in the Phanerozoic*, O. H. Walliser, Ed. (Springer-Verlag, Berlin, 1996), pp. 35–51.
6. A gradually more destabilized Frasnian marine ecosystem by series of comet showers is hypothesized in (9), and the Frasnian-Famennian mass extinction is suggested in (10) to have been triggered by a rapid global cooling that followed Alamo impact-produced anomalous greenhouse interval.
7. A. Hallam, P. B. Wignall, *Mass Extinctions and Their Aftermath* (Oxford Univ. Press, Oxford, 1997).
8. The subsurface Woodleigh impact structure in Western Australia (of disputed size; diameter estimates range from around 40 km to 120 km) was first linked to the Permian-Triassic mass extinction and later to the Late Devonian mass extinction [see (11)], but neither time of impact nor its magnitude seem yet to be well constrained (12).
9. C. A. Sandberg *et al.*, *Geol. Soc. Am. Spec. Pap.* **356**, 473 (2002).
10. G. R. McGhee Jr., *Palaeogeogr. Palaeoclimatol. Palaeoecol.* **176**, 47 (2001).
11. S. D. Uysal *et al.*, *Earth Planet. Sci. Lett.* **192**, 218 (2001).
12. P. R. Renne *et al.*, *Earth Planet. Sci. Lett.* **201**, 247 (2002).




Fig. 1. Two selected impact events plotted against percent extinctions of "well-preserved" marine genera through 16 substages of the Devonian [adapted with permission from figure 6 of Sepkoski (5); copyright © 1996 by Springer-Verlag]. As in the original diagram, end-stage extinction data are not plotted at the graphed stage boundary lines, but slightly before them. Loch, Lochkovian; Prag, Pragian; Emsi, Emsian.

11 August 2003; accepted 2 December 2003