



Comment on "Impacts of Fine Root Turnover on Forest NPP and Soil C Sequestration Potential"

Yiqi Luo, *et al.*

Science **304**, 1745c (2004);

DOI: 10.1126/science.1098080

The following resources related to this article are available online at www.sciencemag.org (this information is current as of May 13, 2008):

Updated information and services, including high-resolution figures, can be found in the online version of this article at:

<http://www.sciencemag.org/cgi/content/full/304/5678/1745c>

A list of selected additional articles on the Science Web sites **related to this article** can be found at:

<http://www.sciencemag.org/cgi/content/full/304/5678/1745c#related-content>

This article **cites 2 articles**, 1 of which can be accessed for free:

<http://www.sciencemag.org/cgi/content/full/304/5678/1745c#otherarticles>

This article appears in the following **subject collections**:

Atmospheric Science

<http://www.sciencemag.org/cgi/collection/atmos>

Technical Comments

http://www.sciencemag.org/cgi/collection/tech_comment

Information about obtaining **reprints** of this article or about obtaining **permission to reproduce this article** in whole or in part can be found at:

<http://www.sciencemag.org/about/permissions.dtl>

Comment on “Impacts of Fine Root Turnover on Forest NPP and Soil C Sequestration Potential”

Norman, OK 73019, USA
and Department of Ecology and
Evolutionary Biology
Fudan University
Shanghai, China

Luther White
Department of Mathematics
University of Oklahoma

Dafeng Hui
Department of Botany and Microbiology
University of Oklahoma

Matamala *et al.* (1) recently highlighted the importance of estimating mean residence time (MRT) of fine root C for understanding soil C dynamics. Using isotopic signals of ^{13}C from two CO_2 experiments as a tracer, they estimated MRT of C through fine roots that ranged from 1.20 to 6.25 years. They obtained these MRT values by fitting an exponential equation to the ^{13}C data with a one-pool model that assumed that newly synthesized C is immediately used for fine root growth.

Actually, however, photosynthetically fixed C is first incorporated into the plant C pool to mix with stored nonstructural carbohydrate (NSC), from which root growth draws C. Thus, the interpretation of isotope data needs to consider NSC storage (2) and to use a two-pool model—that is, one that includes both NSC and fine roots. We developed such a model (3) and estimated MRT of fine roots from the isotope data presented by Matamala *et al.* The resulting MRT values that we derive from this model are 2.37 years for pine roots of <1 mm, 2.01 years for pine

roots of 1 to 2 mm, and 6.06 years for pine roots of 2 to 5 mm (4), shorter by 44.5%, 64.8%, and 3.0%, respectively, than the estimates in (1). For sweetgum roots of <1 mm and 1 to 2 mm, the two-pool model gives estimated MRTs of 1.19 and 1.32 years, respectively—0.8% and 56.0% less than the Matamala *et al.* estimates (1).

Whereas the sums of MRT of fine roots and plant NSC estimated from our two-pool model are similar to the estimates from the one-pool model (1), separating plant NSC from fine root MRT will lead to estimates of soil C sequestration that are different from those of Matamala *et al.* (1). In addition, data points immediately after CO_2 fumigation, which are absent for approximately 10 months of that study, are particularly important for estimation of MRT of plant NSC and need to be collected in future studies.

Yiqi Luo*

Department of Botany and Microbiology
University of Oklahoma

*To whom correspondence should be addressed. E-mail: yluo@ou.edu

References and Notes

1. R. Matamala, M. A. González-Meler, J. D. Jastrow, R. J. Norby, W. H. Schlesinger, *Science* **302**, 1385 (2003).
2. Y. Luo, *Global Change Biol.* **9**, 1118 (2003).
3. The two-pool model is $F(t) = \frac{\beta e^{-\alpha t} - \alpha e^{-\beta t}}{\beta - \alpha}$, where $F(t)$ is old C remaining at time t , and α and β are mean residence times of NSC in plant and fine roots, respectively (5).
4. When we used data from the root ingrowth cores placed 10 months after the initiation of the CO_2 experiments and harvested 24 months after the initiation of the CO_2 experiments to constrain the estimation of MRT of plant NSC, the estimated MRT of fine roots with the two-pool model was still substantially smaller than the value using the one-pool model.
5. Y. Luo, L. White, unpublished data.

8 March 2004; accepted 26 May 2004