

The following resources related to this article are available online at www.sciencemag.org (this information is current as of November 14, 2009):

Updated information and services, including high-resolution figures, can be found in the online version of this article at:

<http://www.sciencemag.org/cgi/content/full/306/5705/2191c>

A list of selected additional articles on the Science Web sites **related to this article** can be found at:

<http://www.sciencemag.org/cgi/content/full/306/5705/2191c#related-content>

This article **cites 18 articles**, 10 of which can be accessed for free:

<http://www.sciencemag.org/cgi/content/full/306/5705/2191c#otherarticles>

This article has been **cited by** 1 articles hosted by HighWire Press; see:

<http://www.sciencemag.org/cgi/content/full/306/5705/2191c#otherarticles>

This article appears in the following **subject collections**:

Technical Comments

http://www.sciencemag.org/cgi/collection/tech_comment

Information about obtaining **reprints** of this article or about obtaining **permission to reproduce this article** in whole or in part can be found at:

<http://www.sciencemag.org/about/permissions.dtl>

Response to Comment on “The Evolution of Modern Eukaryotic Phytoplankton”

Falkowski *et al.* (1) examined when, why, and how a diverse group of eukaryotic phytoplankton, which overwhelmingly contain red plastids, rose to ecological prominence in Mesozoic time and continue to dominate the contemporary oceans. Our analysis included the fossil record of thecate dinoflagellates, coccolithophores, and diatoms; biochemical composition of extant taxa and their phylogenetic relationships; geochemical reconstructions of ocean paleochemistry; eustatic changes in sea level; and ecosystem interactions. One facet in our analysis was the portable plastid hypothesis (2), which was included to accommodate the observation that several distinct clades of eukaryotic algae (e.g., heterokonts, haptophytes, cryptophytes, and dinoflagellates) contain secondary plastids derived from a common ancestral red alga.

A comparative analysis of plastid genomes led Grzebyk *et al.* (2) to conclude that not only are more genes retained in red plastids than in green plastids but also many of the retained genes play critical roles in photosynthetic electron transport and carbon fixation. For example, genes that encode key components of both photosystems, ferredoxin, the ATP synthase and, perhaps most important, the small subunit of ribulose 1,5-bisphosphate carboxylase/oxygenase are present in extant red, but not green, plastids. Although originally based on a small number of plastid genomes, the hypothesis subsequently has been supported by analyses of other chromophytes (3, 4). Based on phylogenetic analyses of nuclear and mitochondrial genomes, the hypothesis assumes that the cells serving as hosts to secondary red plastids do not share a recent common ancestor and, hence, that the plastids in each clade were obtained from independent endosymbiotic events (5, 6). As such, the portable plastid hypothesis implicitly conflicts with the “chromalveolate hypothesis” (7), which proposes that all algae believed to possess secondary red plastids acquired them by a single common endosymbiosis.

There is some evidence that secondary red plastids are derived from different taxa within the red algae (8, 9). However, most analyses that consider the phylogeny of plastid-encoded genes, as well as plastid-targeted genes encoded in the nucleus, indicate that

all secondary red plastids are derived from a common ancestral algal clade (10–12). In contrast, however, phylogenetic analyses of nuclear-encoded genes that are not plastid targeted (e.g., 18S rRNA and cytosolic GAPDH) do not support the proposition that cryptophytes, haptophytes, heterokonts, and alveolates (including dinoflagellates) recently diverged from a common ancestor (5, 13–16). Similarly, mitochondrial genome analyses and ultrastructural features do not comport with a recent common ancestor of chromophyte host cells (17, 18). Paradoxically, there is evidence that the ancestor of heterokonts and alveolates was not photosynthetic. Thus, while the chromalveolate hypothesis is consistent with current phylogenetic data that support a common origin of all secondary red plastids, it is not strongly supported by phylogenetic analyses of the host cells.

A major problem with the chromalveolate hypothesis is that it requires multiple plastid losses in the evolution of alveolates and heterokonts. If the basal groups of alveolates and heterokonts contained a plastid, how and why did nonphotosynthetic alveolates, such as Ciliates, Colpodellids, and Perkinsids (19), and several basal groups of dinoflagellates, lose their plastids? Plastid losses must also be invoked to account for basal heterotrophic heterokonts (20). Plastid losses are not explained by the chromalveolate hypothesis but are not required by the portable plastid hypothesis. Hence, while the chromalveolate hypothesis aims at making a single red plastid acquisition the most parsimonious event in the evolution of secondary symbionts, it is hardly the most parsimonious hypothesis.

Plastid portability is not limited to secondary endosymbiosis of red plastids, but also holds for tertiary endosymbioses and kleptoplastidy. Tertiary red plastids were acquired on at least three occasions in dinoflagellates: from cryptophytes, diatoms, or haptophytes. There are no known tertiary green plastids. Kleptoplastidy (the capability for heterotrophic organisms to temporarily retain functional photosynthetic plastids from algal prey) occurs in dinoflagellates, ciliates, foraminifera, and mollusks. The vast majority of the retained plastids are obtained from chromophytes (21, 22).

Although the vast majority of plastid-targeted genes present in the primary algal cell nucleus were clearly transferred to the secondary host as part of the endosymbiotic process, secondary plastid associations are relatively rare. Assuming that all secondary red plastids originated from a single endosymbiotic event (as the chromalveolate hypothesis proposes) would give even more support to this claim. Although Keeling *et al.* (23) claim that red plastids are no more portable than green plastids, secondary green plastid-containing algae are rare in the contemporary oceans. If green plastids were as portable as red plastids, why is the eukaryotic phytoplankton community in the contemporary ocean dominated by such a diverse group of secondary symbionts that contain red plastids?

Although we believe that current genomic data support the portable plastid hypothesis, the explosion of genomic information expected in the next several years will provide the opportunity to test this hypothesis and the competing chromalveolate alternative.

Daniel Grzebyk

*Environmental Biophysics and
Molecular Ecology Program
Institute of Marine and Coastal Sciences
Rutgers, The State University of New Jersey
71 Dudley Road
New Brunswick, NJ 08901, USA*

Miriam E. Katz

*Department of Geological Sciences
Rutgers, The State University of New Jersey
Piscataway, NJ 08854, USA*

Andrew H. Knoll

*Department of Organismal and
Evolutionary Biology
Harvard University
Cambridge, MA 02138, USA*

Antonietta Quigg*

*Environmental Biophysics and
Molecular Ecology Program
Institute of Marine and Coastal Sciences
Rutgers, The State University of New Jersey
New Brunswick, NJ*

John A. Raven

*Division of Environmental and
Applied Biology
University of Dundee at SCRI
Scottish Crop Research Institute
Invergowrie, Dundee DD2 5DA, UK*

Oscar Schofield

*Environmental Biophysics and
Molecular Ecology Program
Institute of Marine and Coastal Sciences
Rutgers, The State University of New Jersey
New Brunswick, NJ*

TECHNICAL COMMENT

F. J. R. Taylor

Department of Earth and Ocean Science
and Department of Botany
University of British Columbia
270 University Boulevard
Vancouver, BC, Canada V6T 1Z4

Paul G. Falkowski†

Environmental Biophysics and
Molecular Ecology Program
Institute of Marine and Coastal Sciences
Rutgers, The State University of New Jersey
New Brunswick, NJ
and

Department of Geological Sciences
Rutgers, The State University of New Jersey
Piscataway, NJ

*Present address: Department of
Marine Biology
Texas A&M University
Galveston, TX 77551, USA

†To whom correspondence should be
addressed.

E-mail: falko@imcs.rutgers.edu

References and Notes

1. P. G. Falkowski et al., *Science* **305**, 354 (2004).
2. D. Grzebyk, O. Schofield, C. Vetrani, P. G. Falkowski, *J. Phycol.* **39**, 259 (2003).
3. J. R. Manhart, personal communication.
4. M. V. Sanchez Puerta, T. R. Bachvaroff, C. F. Delwiche, *58th Annual Meeting of the Phycological Society of America, Williamsburg VA, 6–12 August 2004*, poster communication (2004).
5. D. Bhattacharya, L. Medlin, *Plant Physiol.* **116**, 9 (1998).
6. C. F. Delwiche, *Am. Nat.* **154**, S164 (1999).
7. T. Cavalier-Smith, *J. Eukaryot. Microbiol.* **46**, 347 (1999).
8. K. M. Müller, M. C. Oliveira, R. G. Sheath, D. Bhattacharya, *Am. J. Bot.* **88**, 1390 (2001).
9. H. S. Yoon, J. D. Hackett, D. Bhattacharya, *Proc. Natl. Acad. Sci. U.S.A.* **99**, 11724 (2002).
10. K.-i. Ishida, B. R. Green, *Proc. Natl. Acad. Sci. U.S.A.* **99**, 9294 (2002).
11. H. S. Yoon, J. D. Hackett, C. Ciniglia, G. Pinto, D. Bhattacharya, *Mol. Biol. Evol.* **21**, 809 (2004).
12. K. Takishita, K.-i. Ishida, T. Maruyama, *Protist* **154**, 443 (2003).
13. N. M. Fast, J. C. Kissinger, D. S. Roos, P. J. Keeling, *Mol. Biol. Evol.* **18**, 418 (2001).
14. J. T. Harper, P. J. Keeling, *Mol. Biol. Evol.* **20**, 1730 (2003).
15. M. E. Katz, Z. V. Finkel, D. Grzebyk, A. H. Knoll, P. G. Falkowski, *Annu. Rev. Ecol. Evol. Syst.* **35**, 523 (2004).
16. K. Takishita, K.-i. Ishida, T. Maruyama, *Protist* **155**, 447 (2004).
17. M. W. Gray et al., *Nucleic Acids Res.* **26**, 865 (1998).
18. M. V. Sanchez Puerta, T. Bachvaroff, C. F. Delwiche, *DNA Res.* **11**, 1 (2004).
19. B. S. Leander, P. J. Keeling, *Trends Ecol. Evol.* **18**, 395 (2003).
20. S. A. Karpov, M. L. Sogin, J. D. Silberman, *Protistology* **2**, 34 (2001).
21. F. J. R. Taylor, D. J. Blackbourn, J. Blackbourn, *J. Fish. Res. Bd. Can.* **28**, 391 (1969).
22. M. Schweikert, M. Elbrächter, *Phycologia* **43**, 614 (2004).
23. P. J. Keeling, J. M. Archibald, N. M. Fast, J. D. Palmer, *Science* **306**, 2191 (2004); www.sciencemag.org/cgi/content/full/306/5705/2191b.
24. Our research is supported by NSF through the Biocomplexity program.

16 September 2004; accepted 29 November 2004
10.1126/science.1105297

Downloaded from www.sciencemag.org on November 14, 2009